

レジメン名称	1コース	催吐性リスク
FP(700/70)+RT(食道癌)	28日間	高度催吐性リスク(day1)

				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
	薬剤	用量	用法																													
	アプレピタント125mg	1Tab	1×化学療法60~90分前	●																												
	アプレピタント80mg	1Tab	1×朝	/	●	●																										
	デキサメタゾン	8mg	1×朝	/	●	●	●																									
滴下順	薬剤	用量	ルート	投与時間	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	21	21	21	21	21	21	
1	生理食塩液	500mL	メイン	2時間	●																											
2	パロノセトロン デキサメタゾン	1バッグ 9.9mg	側管	30分	●																											
3	硫酸マグネシウム 生理食塩液	8mEq 50mL	側管	30分	●																											
4	フルオロウラシル 生理食塩液	700mg/m <sup>2</sup> 1300mL	メイン	24時間	●	●	●	●																								
5	シスプラチン 生理食塩液	70mg/m <sup>2</sup> 350mL	側管	2時間	●																											
6	20%マンニトール	300mg	側管	30分	●																											
7	ソリタT3	500mL	側管	2時間	●																											
8	ソリタT3	500mL	側管	2時間	●																											